



Contents lists available at ScienceDirect

Chinese Journal of Chemical Engineering

journal homepage: www.elsevier.com/locate/CJChE

Separation Science and Engineering

Orthogonal array design for the optimization of stripping Sr(II) from ionic liquids using supercritical CO₂☆

Tao Yang, Jing Fu, Qingde Chen, Xinghai Shen*

Beijing National Laboratory for Molecular Sciences (BNLMS), Fundamental Science on Radiochemistry and Radiation Chemistry Laboratory, College of Chemistry and Molecular Engineering, Peking University, Beijing 100871, China

ARTICLE INFO

Article history:

Received 20 March 2016

Received in revised form 5 May 2016

Accepted 9 May 2016

Available online xxxxx

Keywords:

Supercritical CO₂ stripping

Ionic liquid

Strontium

Orthogonal array design

ABSTRACT

The strontium ions extracted from the aqueous phase into 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide (C₂mimNTf₂) with dicyclohexyl-18-crown-6 (DCH18C6) was stripped effectively by supercritical CO₂ (sc-CO₂). Hexafluoroacetylacetone (HFAA)–acetonitrile was found to be an excellent modifier of sc-CO₂ to enhance the stripping efficiency. In the orthogonal array design (OAD), OA₂₅ (5⁵) matrix was employed to optimize the stripping of Sr(II) from the DCH18C6–C₂mimNTf₂ system. Effects of five experimental factors: temperature, pressure, concentration of HFAA, static and dynamic extraction times as well as each factor at five-levels on the stripping of Sr(II) were optimized. The effects of these parameters were treated by the analysis of variance (ANOVA). The results showed that Sr(II) could be nearly 100% extracted from the IL phase at 308 K, 30 MPa, 40 min of dynamic extraction and 60 mmol·L⁻¹ HFAA in acetonitrile, respectively. Finally, the stripping mechanism was studied by ESI-MS.

© 2016 The Chemical Industry and Engineering Society of China, and Chemical Industry Press. All rights reserved.

1. Introduction

The long lifetime strontium ion is one of the main high heat load, highly radioactive fission products in high level liquid wastes (HLLW) during the reprocessing of spent nuclear fuel. Many advantages were shown to remove Sr(II) from HLLW before the final geological disposal [1]. So far, solvent extraction has been a main method of separating Sr(II) from HLLW, and the ionic liquids (ILs), holding numerous excellent properties and becoming environmentally benign “green solvents” alternative to volatile organic solvents [2–7], have been employed in the extraction of Sr(II).

Dai *et al.* [8] first used ILs as solvents and DCH18C6 as extractant to extract Sr(II) from the aqueous phase. The distribution coefficient (*D*) reached 1.1×10^4 , which was much higher than those of the conventional extraction systems. For example, the *D* values of the extraction systems with C₆H₅CH₃, CHCl₃, 2-octanol and 1,1,2,2-tetrachloroethane as diluents were only 0.76 [8], 0.77 [8], 6.5 [9] and 10.8 [10], respectively. Dietz and Dzielawa [11] proposed that the extraction of Sr(II) by IL systems was based on a cation exchange mechanism. Our research group studied the effect of the structure of ionic liquid, the acidity and the addition of inorganic salt in the aqueous phase on the extraction efficiency of Sr(II) by the DCH18C6–C_{*n*}mimNTf₂ (*n* = 2, 4, 6) system, and further confirmed the cation exchange mechanism [1]. In addition,

we found that the γ -radiation stabilities of ILs were excellent during the extraction of Sr(II) with DCH18C6 [12,13], which is conducive to fit to the high irradiation environment in the spent nuclear fuel reprocessing.

Although ILs have lots of advantages towards the extraction of metal ions, the stripping, which is as important as the extraction process in the application, still remains a challenge. Our research group tried to study the stripping of Sr(II) with K₂SO₄ and the stripping efficiency reached 99% [1]. But there are two serious obstacles: cross contamination between two phases and loss of ILs. Electrodeposition is another method to recover metal ions from the IL phase [14,15]. However, this method applies at many strict conditions, *i.e.*, only for highly electropositive elements, and requiring the electrochemical window of an IL matching the reduction potential of a metal ion.

Supercritical CO₂ (sc-CO₂) is considered as another “green solvent”, because CO₂ can be recycled and generates no wastes. Sc-CO₂ has many advantages including enhanced diffusivity, chemical inertness, non-flammability and low cost [16–18]. Sc-CO₂ has been employed to extract various metal ions such as lanthanides, actinides, alkali metals from aqueous or solid matrices [19–32]. Due to the limited mutual solubility between sc-CO₂ and IL, that is, the insolubility of IL in sc-CO₂ and the solubility of sc-CO₂ in IL [33–35], a feasible method to recover metal ions from IL phase by sc-CO₂ has been put forward. Some metal ions were stripped effectively by sc-CO₂ from IL phase [36–40]. However, the mechanism of stripping by sc-CO₂ has been rarely reported. Our research group recently used ESI-MS to study the mechanism of stripping U(VI) from the CMPO–C₂mimNTf₂ system by TOPO-modified sc-CO₂ [40].

☆ Supported by the National Natural Science Foundation of China (91226112).

* Corresponding author. Tel.: +86 10 62765915; Fax: +86 10 62759191.

E-mail address: xshen@pku.edu.cn (X. Shen).

To the best of our knowledge, although DCH18C6 as the extractant in $C_2mimNTf_2$ performs excellently for the extraction of Sr(II), there have been no effective methods so far to strip it from the DCH18C6– $C_2mimNTf_2$ system with high stripping efficiency, no cross-contamination, and no loss of ILs.

In this study, we aim to use sc- CO_2 to strip Sr(II) from the DCH18C6– $C_2mimNTf_2$ system. Due to the strong polarity of $DCH18C6 \cdot Sr^{2+}$ but nonpolarity of CO_2 , the solubility of crown ether complexes in sc- CO_2 is very small. It is well known that fluorinated metal chelates are CO_2 -philic [41], thus utilizing fluorinated anion to combine crown ether-metal cation as ion-pair is a favorable method. Wai *et al.* [23] successfully extracted Sr(II) from aqueous media into sc- CO_2 employing DCH18C6 and a fluorinated anion, *i.e.*, $CF_3(CF_2)_6CO_2^-$ (PFOA⁻) or $CF_3(CF_2)_6CF_2SO_3^-$ (PFOSA⁻). Another method to improve the solubility of crown ether complexes is adding cosolvent to sc- CO_2 , such as acetonitrile, methanol, and acetone to enhance its polarity. Considering the two aspects in favor of the solubility of crown ether complexes in sc- CO_2 , we chose a fluorinated metal chelate hexafluoroacetylacetone (HFAA) diluted in acetonitrile as the modifier.

Meanwhile, the influences of temperature, pressure, static stripping time, dynamic stripping time, and the concentration of HFAA on the stripping efficiency of Sr(II) by sc- CO_2 were investigated systematically. Due to the difficulty in selecting the optimized conditions, the orthogonal array design (OAD) as an artful method was used to evaluate objectively the whole conditions in the sc- CO_2 stripping progress. The results of OAD experiments were investigated by the analysis of variance (ANOVA) [42,43].

2. Experimental

2.1. Materials

$C_2mimNTf_2$ was synthesized and purified according to the reported methods in the literature [44–46]. The purity was above 99% as determined by elemental analyses and 1H NMR. Dicyclohexano-18-crown-6 (DCH18C6) was received from INET, Tsinghua University and purified by recrystallization in *n*-heptane. The purity of final product was above 98%. 2,2-Dimethyl-6,6,7,7,8,8,8-heptafluoro-3,5-octanedione (HFOD, 99%, J&K), hexafluoroacetylacetone (HFAA, 99%, J&K), thenoyltrifluoroacetone (HTTA, 99%, J&K) and perfluorooctanoic acid (HPFOA, 98%, J&K) were used as received. The chemical structures of $C_2mimNTf_2$, DCH18C6, HFAA, HTTA, HPFOA and HFOD are shown in Fig. 1. $Sr(NO_3)_2$ and other reagents were of analytical grade. CO_2 (99.95%) from Reijiye in China was used in all experiments.

2.2. Instruments

All stripping experiments were performed in a lab supercritical fluid extraction apparatus (100 ml, SEPAREX, France). The concentration of Sr(II) was analyzed by an Inductively Coupled Plasma-Atomic Emission Spectrometer (ICP-AES, Leeman, USA). The high-resolution mass spectrometry used with electrospray ionization (ESI-HRMS) was done on a Fourier transform ion cyclotron resonance mass spectrometer, APEX IV (Bruker, USA). Microwave digestion was completed in the instrument of CEM MARS6 (USA).

2.3. Methods

The stripping samples were obtained after extracting Sr(II) from the aqueous phase by DCH18C6– $C_2mimNTf_2$. The concentration of Sr(II) was about $0.025 \text{ mol} \cdot L^{-1}$ in the $C_2mimNTf_2$ phase. $0.35 \text{ ml } C_2mimNTf_2$ solution containing $0.025 \text{ mol} \cdot L^{-1}$ of strontium was added in a 10 ml glass-basket, which was fixed in the high-pressure extraction vessel with stirring. The stripping process was allowed to take place under static and dynamic extraction conditions. Under static extraction condition, 5 ml modifier was injected in vessel and kept for some time. The




Fig. 1. Molecular structures of $C_2mimNTf_2$ (a), DCH18C6 (b), HFAA (c), HFOD (d), HTTA (e), and HPFOA (f).

flow rate of CO_2 was fixed at $50 \text{ g} \cdot \text{min}^{-1}$, and the modifier was the $0.5 \text{ ml} \cdot \text{min}^{-1}$ of HFAA–acetonitrile solution under dynamic extraction condition.

When the stripping progress was completed, the system was allowed to slowly depressurize and the IL sample was then removed from the cell and analyzed. The stripping efficiency was calculated based on the contents of Sr(II) in the initial and the residual IL phases, which were analyzed by ICP-AES after microwave digestion. The mass spectra of the stripping product and the IL phases were measured with ESI-MS.

3. Results and Discussion

3.1. Experimental design and data analysis

As known, the density of sc- CO_2 decreases with raising the temperature and reducing the pressure. Generally, the solubility of the solids in the sc- CO_2 mainly depends on the density of the solvent. However, the extraction efficiency does not only depend on the density of sc- CO_2 [43,47]. For example, the rising of temperature in the course of stripping could increase the volatility of the analytes, accelerate the complexation reactions and desorption of the analytes from the IL phase, which increased the extraction efficiency [43,47]. With respect to the pressure, the interaction between sc- CO_2 and IL was reduced because of the decrease of the diffusion coefficient at a higher density, which can reduce the extraction efficiency [43,47]. Therefore, the temperature (T) and pressure (P) should be optimized. In addition, static (t_s) and dynamic (t_d) extraction times as well as the concentration of HFAA (C) are also important factors on the stripping efficiency. Thus, a five-factor, five-level OAD with an $OA_{25} (5^5)$ matrix was designed to evaluate the effects of the following factors: T , P , t_s , t_d and C . In order to obtain the best condition for the stripping of Sr(II), 25 experiments were performed. The factors and their corresponding levels as well as the results of the stripping experiments are reported in Table 1.

Table 1
Results of $OA_{25}(5^5)$ experimental design for the extraction of Sr(II)

Trial number	T/K	P/MPa	t_s /min	t_d /min	C/mmol·L ⁻¹	E/%
1	308	10	15	10	60	22 ± 3
2	308	15	20	20	80	43 ± 2
3	308	20	25	30	100	64 ± 3
4	308	25	30	40	120	97 ± 4
5	308	30	35	50	140	96 ± 6
6	318	10	25	20	120	9 ± 2
7	318	15	30	30	140	50 ± 6
8	318	20	35	40	60	91 ± 1
9	318	25	15	50	80	96 ± 1
10	318	30	20	10	100	70 ± 6
11	328	10	35	30	80	3 ± 1
12	328	15	15	40	100	46 ± 1
13	328	20	20	50	120	92 ± 4
14	328	25	25	10	140	52 ± 2
15	328	30	30	20	60	57 ± 3
16	338	10	30	50	100	4 ± 5
17	338	15	35	10	120	10 ± 1
18	338	20	15	20	140	51 ± 1
19	338	25	20	30	60	61 ± 3
20	338	30	25	40	80	93 ± 1
21	348	10	20	40	140	6 ± 2
22	348	15	25	50	60	4 ± 4
23	348	20	30	10	80	18 ± 3
24	348	25	35	20	100	28 ± 3
25	348	30	15	30	120	85 ± 1

The mean value of each stripping efficiency for the respective factors and at each level was calculated according to the assignment of the experiment (Table 1). For example, the stripping efficiency of the five trials at 308 K was evaluated as mean value of the corresponding five runs (Trial number 1–5). The mean values of the five-levels of each factor reveal how the stripping efficiency changes when changing the level of that factor. Fig. 2 shows stripping efficiency as a function of levels of the studied factors. In all instances, it should be noted that each calculated stripping efficiency is the average of five measurements, and in each of which the interested parameter was kept constant, and all the other parameters were changed (Table 1).

The ANOVA results for calculated models are shown in Table 2. The ANOVA indicates that the temperature and pressure of sc-CO₂ play key roles and the dynamic time plays important role in the stripping of Sr(II) from the IL phase, whereas, in the selected range the static time and the concentration of HFAA do not have significant effects on the stripping of Sr(II).

3.2. Effect of extraction temperature

The variation of temperature during stripping affects the density of sc-CO₂, the volatility of the analytes and desorption of the analytes from the IL phase. At higher temperatures, the density of sc-CO₂

Table 2
ANOVA of the Sr(II) extractions

Source of variance	Sum of square	Degree of freedom	Variance	F	F_{α}	Significant level
T	4457	4	1114	5.8	$F_{0.05} = 3.26$	**
P	17257	4	4314	22.4	$F_{0.01} = 5.41$	**
t_s	988	(4)	247	1.3		
t_d	3730	4	933	4.8		*
C	689	(4)	172	0.9		
Pooled error	2310	12	193			

Note: $F = \text{Variance (factor)} / \text{Variance (error)}$.

F_{α} is obtained from F value table for analysis of variance.

** stands for playing key role, * stands for playing important role.

decreases and analytes become more volatile leading to lower efficiency as shown in Fig. 2(a).

The ANOVA of the results shows that the temperature plays a key role in the extraction of Sr(II). Therefore, we ought to select a lower extraction temperature (308 K) as the optimum temperature.

3.3. Effect of extraction pressure

Solubility of a solute in sc-CO₂ depends on a balance between sc-CO₂ density and solute vapor pressure, both of which are controlled by the pressure of sc-CO₂. As can be learnt from the ANOVA of the results, the pressure of the sc-CO₂ also plays a key role in the stripping of Sr(II) from the IL phase. This means that the stripping efficiency is enhanced by an increase in the pressure. As shown in Fig. 2(b), the optimized pressure is 30 MPa.

3.4. Effect of static and dynamic extraction times

In order to achieve high stripping efficiency, the static extraction can make the penetration of sc-CO₂ in the IL phase better than the dynamic extraction. The dynamic extraction follows static extraction to enhance the solubility of analyte in sc-CO₂.

Based on the ANOVA calculations, the dynamic time plays an important role in the stripping of Sr(II) from the IL phase. According to Fig. 2(d), when the dynamic time is increased, the stripping efficiency of Sr(II) firstly increases and then decreases, and the best dynamic time is 40 min. On the contrary, the static time has no significant effects on the stripping efficiency. For a rapid extraction of Sr(II) from the IL phase and on the basis of economy principle, one can select a shorter static time (15 min).

3.5. Effect of the concentration of HFAA

Based on the ANOVA calculations, the concentration of HFAA has no significant effects on the stripping of Sr(II) from the IL phase in the selected range (60–140 mmol·L⁻¹). Therefore, we select 60 mmol·L⁻¹




Fig. 2. Effect of temperature (a), pressure (b), static (c) and dynamic (d) extraction times and the concentration of HFAA (e) on the stripping of Sr(II) from DCH18C6–C₂mimNTf₂.

as the optimum concentration of HFAA according to the economy principle.

On the basis of the above results, the optimum values of the selected factors (temperature, pressure, static and dynamic extraction times and concentration of HFAA) for stripping of Sr(II) are 308 K, 30 MPa, 15 min, 40 min and 60 mmol·L⁻¹, respectively. Further experiment was performed under the proposed conditions and the stripping efficiency was about 100%.

In order to obtain the better conditions, we further optimized static stripping time and concentration of HFAA. When the static time was 15, 10, 5 and 0 min respectively at 308 K, 30 MPa, 40 min of dynamic stripping time and 60 mmol·L⁻¹ HFAA in acetonitrile, the stripping efficiency of Sr(II) always reached 100%. The effect of the concentration of HFAA is shown in Fig. 3. The stripping efficiency is reduced when the concentration of HFAA decreases. So we can obtain the optimized conditions, namely, no static stripping time and the concentration of HFAA is 60 mmol·L⁻¹ at 308 K, 30 MPa, 40 min of dynamic stripping time.




Fig. 3. Effect of concentration of HFAA on the stripping efficiency of Sr(II) at 308 K, 30 MPa, 40 min of dynamic stripping time from C₂mimNTf₂.

3.6. Stripping mechanism

The above results demonstrate that Sr(II) can be effectively stripped from the IL phase into the sc-CO₂ phase with HFAA–acetonitrile solution as the modifier. Acetonitrile as a co-solvent can enhance the polarity of CO₂ to increase the solubility of the complexes of Sr(II) in the modified sc-CO₂.

The stripping efficiency of Sr(II) from the IL phase can reach 100% and less than 5% of IL is lost during the stripping experiment under the optimal condition: 308 K, 30 MPa, 40 min of dynamic stripping time and 60 mmol·L⁻¹ HFAA in acetonitrile.

We investigated the original IL phase, the recovered IL phase, and the stripping product by ESI-MS (Fig. 4(a-c)), respectively. The compositions of the fragments and their corresponding complexes are listed in Table 3.

Cationic complexes [C₂mim-DCH18C6-NTf₂-Sr-2H₂O]²⁺, [Sr-NO₃-H₂O]⁺, [Sr-NO₃-FAA-Sr]²⁺, [DCH18C6-Sr-FAA]⁺, [C₂mim-NTf₂-DCH18C6-FAA-Sr]⁺ and [DCH18C6-Sr-NTf₂]⁺ are observed in the ESI spectrum of the stripping product (Fig. 4(c)), which indicates that various kinds of the complexes of Sr(II) were formed in the stripping process. FAA⁻ not only competes with DCH18C6 to associate with Sr²⁺, but also strips Sr²⁺ with DCH18C6 synergistically. According to the above results, one can infer that the affinity of Sr²⁺ with FAA⁻ is larger than




Fig. 4. Positive ESI spectra of the original IL phase (a), the recovered IL phase (b), and the stripping product (c).

that with DCH18C6, and the complexes of Sr(II) associating with FAA⁻ become more soluble in sc-CO₂. Therefore, with FAA⁻ as the modifier, sc-CO₂ can strip strontium from the DCH18C6–C₂mimNTf₂ system effectively. With respect to the recovery of Sr(II) from the chelate, two methods could be applied. If the recovery of complexing agents is not necessary, Sr(NO₃)₂ aqueous solution could be obtained via the digestion of the stripping product. If the recovery of ligands is obligatory, SrSO₄ precipitate could be collected by washing the organic solution of the stripping product with the aqueous solution containing SO₄²⁻.

The cationic complexes [DCH18C6-Na]⁺ and [DCH18C6-NH₄]⁺ demonstrate the existence of the extractant DCH18C6 in the stripping product and illustrate that part of DCH18C6 can be brought out by modified sc-CO₂. The cationic complex containing [C₂mim-NTf₂-C₂mim]⁺ is also observed in the stripping product (Fig. 4(c)), which means that the IL was brought out partly by sc-CO₂ with HFAA–acetonitrile as the modifier. According to the literature, the dissolving capacity of ILs in sc-CO₂ can be enhanced significantly, when the sc-CO₂ phase contains polar organic solvents acting as co-solvents [35]. The effect of a co-solvent on the solubility of sc-CO₂ for the ILs primarily depends on its polarity and concentration in sc-CO₂ [48,49]. Therefore, we believe that the loss of the ILs can be avoided to some extent if the stripping conditions are further optimized.

Table 3

The compositions of the fragments and their corresponding complexes of the original IL phase, the recovered IL phase and stripping product

<i>m/z</i>	Cationic complex
<i>The original IL phase</i>	
167.4	$[\text{Sr-NO}_3\text{-H}_2\text{O}]^+$
251.1	$[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim-H}]^{2+}$
390.3	$[\text{DCH18C6-NH}_4]^+$
411.2	$[\text{DCH18C6-K}]^+$
443.3	$[\text{C}_2\text{mim-NTf}_2\text{-DCH18C6-Sr-2H}_2\text{O}]^{2+}$
502.1	$[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim}]^+$
740.1	$[\text{DCH18C6-Sr-NTf}_2]^+$
773.5	$[\text{2C}_2\text{mim-2NTf}_2\text{-2DCH18C6-H}_2\text{O-2H}]^{2+}$
<i>The recovered IL phase</i>	
167.4	$[\text{Sr-NO}_3\text{-H}_2\text{O}]^+$
251.1	$[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim-H}]^{2+}$
411.2	$[\text{DCH18C6-K}]^+$
502.1	$[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim}]^+$
773.5	$[\text{2C}_2\text{mim-2NTf}_2\text{-2DCH18C6-H}_2\text{O-2H}]^{2+}$
<i>The stripping product</i>	
167.4	$[\text{Sr-NO}_3\text{-H}_2\text{O}]^+$
222.4	$[\text{Sr-NO}_3\text{-FAA-Sr}]^{2+}$
390.3	$[\text{DCH18C6-NH}_4]^+$
395.2	$[\text{DCH18C6-Na}]^+$
443.3	$[\text{C}_2\text{mim-NTf}_2\text{-DCH18C6-Sr-2H}_2\text{O}]^{2+}$
502.1	$[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim}]^+$
667.1	$[\text{DCH18C6-Sr-FAA}]^+$
740.1	$[\text{DCH18C6-Sr-NTf}_2]^+$
1058.2	$[\text{C}_2\text{mim-NTf}_2\text{-DCH18C6-FAA-Sr}]^+$

In the ESI spectrum of the recovered IL phase (Fig. 4(b)), the signals of the cationic complexes $[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim}]^+$ and $[\text{C}_2\text{mim-NTf}_2\text{-C}_2\text{mim-H}]^{2+}$ are observed, which are consistent with those in the ESI spectrum of the original IL phase (Fig. 4(a)). And the cationic complexes $[\text{Sr-NO}_3\text{-H}_2\text{O}]^+$ exist in the recovered IL phase, but the intensity is very weak. This indicates that Sr(II) has been stripped almost completely, which is in agreement with the results of ICP-AES. Thus, highly efficient stripping of Sr(II) from the DCH18C6–C₂mimNTf₂ system was achieved in this work.

Combined with our previous work about the extraction of Sr(II) by DCH18C6–C₂mimNTf₂ [1], a complete cycle including extraction and stripping processes for Sr(II) with both IL and sc-CO₂ can be established. The whole process is illustrated in Fig. 5.

Finally, we evaluated the other three fluorinated metal chelates HTTA, HPOD, and HFPOA, respectively, diluted in acetonitrile at the optimized conditions of HFAA, and the stripping efficiencies are 66%, 87%




Fig. 6. Effect of different modifiers diluted in acetonitrile on the stripping of Sr(II) from DCH18C6–C₂mimNTf₂.

and 91% (Fig. 6). They can also be chosen as modifiers and their stripping efficiencies could be further optimized by OAD.

4. Conclusions

The stripping of Sr(II) from the DCH18C6–C₂mimNTf₂ system using the modified sc-CO₂ was studied. It was found that HFAA–acetonitrile is an efficient modifier to facilitate the stripping of Sr(II) from the IL phase. By the method of OAD and ANOVA, nearly 100% of Sr(II) could be successfully stripped off under the following conditions: 308 K, 30 MPa, 40 min of dynamic stripping time and 60 mmol·L⁻¹ HFAA in acetonitrile.

The present work demonstrates the feasibility of employing sc-CO₂ to strip and promises a highly efficient extraction–stripping cycle for IL-based Sr(II) recovery process.

Nomenclature

<i>C</i>	concentration of HFAA, mmol·L ⁻¹
<i>P</i>	pressure, MPa
<i>T</i>	temperature, K
<i>t_d</i>	dynamic extraction time, min
<i>t_s</i>	static extraction time, min

Subscripts

<i>d</i>	dynamic extraction
<i>s</i>	static extraction




Fig. 5. A diagram showing the extraction–stripping procedure for strontium.

References

- [1] C. Xu, X.H. Shen, Q.D. Chen, H.C. Gao, Investigation on the extraction of strontium ions from aqueous phase using crown ether-ionic liquid systems, *Sci. China Ser. B Chem.* 52 (2009) 1858–1864.
- [2] H.W. Liu, T. Yang, Q.D. Chen, X.H. Shen, Extraction behaviors of ionic liquid systems and application perspectives in reprocessing of spent nuclear fuel, *J. Nucl. Radiochem.* 37 (2015) 286–309.
- [3] X.Q. Sun, H.M. Luo, S. Dai, Ionic liquids-based extraction: A promising strategy for the advanced nuclear fuel cycle, *Chem. Rev.* 112 (2012) 2100–2128.
- [4] R. Hagiwara, Y. Ito, Room temperature ionic liquids of alkylimidazolium cations and fluoroanions, *J. Fluor. Chem.* 105 (2000) 221–227.
- [5] R.D. Rogers, K.R. Seddon, Ionic liquids – solvents of the future? *Science* 302 (2003) 792–793.
- [6] H. Zhao, S.Q. Xia, P.S. Ma, Use of ionic liquids as 'green' solvents for extractions, *J. Chem. Technol. Biotechnol.* 80 (2005) 1089–1096.
- [7] C. Chiappe, D. Pieraccini, Ionic liquids: Solvent properties and organic reactivity, *J. Phys. Org. Chem.* 18 (2005) 275–297.
- [8] S. Dai, Y.H. Ju, C.E. Barnes, Solvent extraction of strontium nitrate by a crown ether using room-temperature ionic liquids, *J. Chem. Soc. Dalton* (1999) 1201–1202.
- [9] L.H. He, X.M. Weng, D.Z. Yang, C.L. Song, Extraction of strontium from high-level active waste with crown-ether III. Extraction of strontium with dicyclohexyl-18-crown-6-octanol-[2], *J. Nucl. Radiochem.* (1994) 18–22.
- [10] Q. Yang, Y.D. Han, D.M. Liu, Extraction of strontium(II) from nitric acid medium with crown-ether, *J. Environ. Sci. Technol.* (1996) 7–9.
- [11] M.L. Dietz, J.A. Dzielawa, Ion-exchange as a mode of cation transfer into room-temperature ionic liquids containing crown ethers: Implications for the 'greenness' of ionic liquids as diluents in liquid-liquid extraction, *Chem. Commun.* (2001) 2124–2125.
- [12] L.Y. Yuan, J. Peng, L. Xu, M.L. Zhai, J.Q. Li, G.S. Wei, Radiation effects on hydrophobic ionic liquid [C₄mim][NTf₂] during extraction of strontium ions, *J. Phys. Chem. B* 113 (2009) 8948–8952.
- [13] L.Y. Yuan, J. Peng, L. Xu, M.L. Zhai, J.Q. Li, G.S. Wei, Influence of gamma-radiation on the ionic liquid [C₄mim][PF₆] during extraction of strontium ions, *Dalton Trans.* (2008) 6358–6360.
- [14] P.Y. Chen, The assessment of removing strontium and cesium cations from aqueous solutions based on the combined methods of ionic liquid extraction and electrodeposition, *Electrochim. Acta* 52 (2007) 5484–5492.
- [15] P. Giridhar, K.A. Venkatesan, S. Subramaniam, T.G. Srinivasan, P.R.V. Rao, Extraction of uranium (VI) by 1.1 M tri-*n*-butylphosphate/ionic liquid and the feasibility of recovery by direct electrodeposition from organic phase, *J. Alloys Compd.* 448 (2008) 104–108.
- [16] C.M. Wai, B. Waller, Dissolution of metal species in supercritical fluids – principles and applications, *Ind. Eng. Chem. Res.* 39 (2000) 4837–4841.
- [17] P. Kumar, A. Pal, M.K. Saxena, K.L. Ramakumar, Supercritical fluid extraction of uranium and thorium from solid matrices, *Desalination* 232 (2008) 71–79.
- [18] W.H. Duan, L.Y. Zhu, S. Jing, Y.J. Zhu, J. Chen, Study on properties of TBP-HNO₃ complex used for direct dissolution of lanthanide and actinide oxides in supercritical fluid CO₂, *Chin. J. Chem.* 25 (2007) 319–322.
- [19] K.E. Laintz, E. Tachikawa, Extraction of lanthanides from acidic solution using tributyl-phosphate modified supercritical carbon-dioxide, *Anal. Chem.* 66 (1994) 2190–2193.
- [20] Y. Meguro, S. Iso, H. Takeishi, Z. Yoshida, Extraction of uranium(VI) in nitric acid solution with supercritical carbon dioxide fluid containing tributylphosphate, *Radiochim. Acta* 75 (1996) 185–191.
- [21] S. Mochizuki, N. Wada, R.L. Smith, H. Inomata, Perfluorocarboxylic acid counter ion enhanced extraction of aqueous alkali metal ions with supercritical carbon dioxide, *Analyst* 124 (1999) 1507–1511.
- [22] S. Mochizuki, N. Wada, R.L. Smith, H. Inomata, Supercritical fluid extraction of alkali metal ions using crown ethers with perfluorocarboxylic acid from aqueous solution, *Anal. Commun.* 36 (1999) 51–52.
- [23] C.M. Wai, Y. Kulyako, H.K. Yak, X.Y. Chen, S.J. Lee, Selective extraction of strontium with supercritical fluid carbon dioxide, *Chem. Commun.* (1999) 2533–2534.
- [24] C.M. Wai, Y.M. Kulyako, B.F. Myasoedov, Supercritical carbon dioxide extraction of caesium from aqueous solutions in the presence of macrocyclic and fluorinated compounds, *Mendeleev Commun.* (1999) 180–181.
- [25] S. Mochizuki, R.L. Smith, H. Inomata, Quantitative extraction of aqueous alkali metal ions using supercritical carbon dioxide and polyethylene glycol ligands, *Chem. Commun.* (2000) 1381–1382.
- [26] Y.H. Lin, C.X. Liu, H. Wu, H.K. Yak, C.M. Wai, Supercritical fluid extraction of toxic heavy metals and uranium from acidic solutions with sulfur-containing organo-phosphorus reagents, *Ind. Eng. Chem. Res.* 42 (2003) 1400–1405.
- [27] S.F. Wang, Y.H. Lin, C.M. Wai, Supercritical fluid extraction of toxic heavy metals from solid and aqueous matrices, *Sep. Sci. Technol.* 38 (2003) 2279–2289.
- [28] T. Shimada, S. Ogumo, K. Sawada, Y. Enokida, I. Yamamoto, Selective extraction of uranium from a mixture of metal or metal oxides by a tri-*n*-butylphosphate. Complex with HNO₃ and H₂O in supercritical CO₂, *Anal. Sci.* 22 (2006) 1387–1391.
- [29] J.S. Wang, K. Chiu, Metal extraction from solid matrices using a two-surfactant microemulsion in neat supercritical carbon dioxide, *Microchim. Acta* 167 (2009) 61–65.
- [30] D.L. Quach, B.J. Mincher, C.M. Wai, Supercritical fluid extraction and separation of uranium from other actinides, *J. Hazard. Mater.* 274 (2014) 360–366.
- [31] A. Rao, P. Kumar, B.S. Tomar, Supercritical fluid extraction of uranium and thorium employing dialkyl amides, *Sep. Purif. Technol.* 134 (2014) 126–131.
- [32] H. Xu, Research of Supercritical Carbon Dioxide Fluid Extraction of Strontium. Master Degree Thesis Institute of China Atom Science Research Institute, Beijing, 2005.
- [33] L.A. Blanchard, D. Hancu, E.J. Beckman, J.F. Brennecke, Green processing using ionic liquids and CO₂, *Nature* 399 (1999) 28–29.
- [34] S.G. Kazarian, B.J. Briscoe, T. Welton, Combining ionic liquids and supercritical fluids: *in situ* ATR-IR study of CO₂ dissolved in two ionic liquids at high pressures, *Chem. Commun.* (2000) 2047–2048.
- [35] S. Keskin, D. Kayrak-Talay, U. Akman, O. Hortacsu, A review of ionic liquids towards supercritical fluid applications, *J. Supercrit. Fluids* 43 (2007) 150–180.
- [36] S. Mekki, C.M. Wai, I. Billard, G. Moutiers, C.H. Yen, J.S. Wang, A. Ouadi, C. Gaillard, P. Hesemann, Cu(II) extraction by supercritical fluid carbon dioxide from a room temperature ionic liquid using fluorinated beta-diketones, *Green Chem.* 7 (2005) 421–423.
- [37] S. Mekki, C.M. Wai, I. Billard, G. Moutiers, J. Burt, B. Yoon, J.S. Wang, C. Gaillard, A. Ouadi, P. Hesemann, Extraction of lanthanides from aqueous solution by using room-temperature ionic liquid and supercritical carbon dioxide in conjunction, *Chem. Eur. J.* 12 (2006) 1760–1766.
- [38] J.S. Wang, C.N. Sheaff, B. Yoon, R.S. Addleman, C.M. Wai, Extraction of uranium from aqueous solutions by using ionic liquid and supercritical carbon dioxide in conjunction, *Chem. Eur. J.* 15 (2009) 4458–4463.
- [39] J. Fu, Q.D. Chen, T.X. Sun, X.H. Shen, Extraction of Th(IV) from aqueous solution by room-temperature ionic liquids and coupled with supercritical carbon dioxide stripping, *Sep. Purif. Technol.* 119 (2013) 66–71.
- [40] J. Fu, Q.D. Chen, X.H. Shen, Stripping of uranium from an ionic liquid medium by TOPO-modified supercritical carbon dioxide, *Sci. China: Chem.* 58 (2015) 545–550.
- [41] K.E. Laintz, J.J. Yu, C.M. Wai, Separation of metal-ions with sodium bis(trifluoroethyl)diithiocarbamate chelation and supercritical fluid chromatography, *Anal. Chem.* 64 (1992) 311–315.
- [42] S.M. Pourmortazavi, S.S. Hajimirsadeghi, I. Kohsari, S.G. Hosseini, Orthogonal array design for the optimization of supercritical carbon dioxide extraction of different metals from a solid matrix with cyanex 301 as a ligand, *J. Chem. Eng. Data* 49 (2004) 1530–1534.
- [43] Y. Yamini, A. Saleh, M. Khajeh, Orthogonal array design for the optimization of supercritical carbon dioxide extraction of platinum(IV) and rhenium(VII) from a solid matrix using cyanex 301, *Sep. Purif. Technol.* 61 (2008) 109–114.
- [44] P. Bonhote, A.P. Dias, N. Papageorgiou, K. Kalyanasundaram, M. Gratzel, Hydrophobic, highly conductive ambient-temperature molten salts, *Inorg. Chem.* 35 (1996) 1168–1178.
- [45] J.G. Huddleston, A.E. Visser, W.M. Reichert, H.D. Willauer, G.A. Broker, R.D. Rogers, Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation, *Green Chem.* 3 (2001) 156–164.
- [46] J.G. Huddleston, H.D. Willauer, R.P. Swatloski, A.E. Visser, R.D. Rogers, Room temperature ionic liquids as novel media for 'clean' liquid-liquid extraction, *Chem. Commun.* (1998) 1765–1766.
- [47] M.Z. Fang, S.Y. Sun, X.D. Zheng, C.J. Lin, Study on the supercritical CO₂ chelating extraction of heavy metals, *Guangdong Chem. Ind.* (2007) 61–63.
- [48] W.Z. Wu, W.J. Li, B.X. Han, T. Jiang, D. Shen, Z.F. Zhang, D.H. Sun, B. Wang, Effect of organic cosolvents on the solubility of ionic liquids in supercritical CO₂, *J. Chem. Eng. Data* 49 (2004) 1597–1601.
- [49] W.Z. Wu, J.M. Zhang, B.X. Han, J.W. Chen, Z.M. Liu, T. Jiang, J. He, W.J. Li, Solubility of room-temperature ionic liquid in supercritical CO₂ with and without organic compounds, *Chem. Commun.* (2003) 1412–1413.